

Supplemental Table S1.

Logic Rules	References
AKT = PIP3	[1]
CCR4 = CCL17	[2]
COX2 = (TNFAR and S1P)	[3][4][5]
EP2 = PGE2	[6]
ERK = MEK	[7][7]
GP130 = IL6	
IKB = not IKK	
IKK = AKT and (not (TNFA and JAK2) or not ROCK or not IRAK1)	[8][9][10]
IL1	
IL13	
IL13R = IL13	
IL1R = IL1	
IL23	
IL23R = IL23	
IL6	
IRAK1 = IL1R	[9]
JAK1 = GP130 or IL13R or TNFA	[8]
JAK2 = GP130 or IL13R or TNFA or IL23R	[11]
MEK = RAF	[12]
MUC1STN = ST6GALNAC1	
P65 = not IKKB	[13][10][14]
PGE2 = COX2	[15]
PI3K = (CCR4 and RAS and not PTEN) or (EP2 and CCR4 and VEGFR and not PTEN)	[16]
PIP3 = PI3K	[16]
PTEN = ROCK and not P65	[16]
RAF = RAS	[12]
RAS = GP130 or EP2 or VEGFR	[17][18]
RHOA = (IL13R and CCR4 and ERK and TGFB)	[2][19][20][21][22]
ROCK = RHOA	[22]
S1P = SPHK1	[23]
SOCS3 = STAT3	[23][24]
SPHK1 = TNFA or ERK	[23][3]
ST6GALNAC1 = ST6GALNAC1gene	
ST6GALNAC1gene = (STAT1 and STAT3 and STAT6), P65	This study
STAT1 = JAK1 or TYK2	[8]
STAT3 = JAK1 or TYK2	[23][25][8]
STAT6 = TYK2 or ROCK	[22][8]
CCL17	
TGFB	
TGFBR = TGFB	
TNFA	
TNFAR = TNFA	
TYK2 = GP130 or IL13R	[2]

VEGF	
VEGFR = VEGF	

References

- [1] A. Carnero, C. Blanco-aparicio, O. Renner, W. Link, and J. F. M. Leal, “The PTEN / PI3K / AKT Signalling Pathway in Cancer , Therapeutic Implications,” pp. 187–198, 2008.
- [2] A. A. Al-Haidari, I. Syk, K. Jirström, and H. Thorlacius, “CCR4 mediates CCL17 (TARC)-induced migration of human colon cancer cells via RhoA/Rho-kinase signaling,” *Int. J. Colorectal Dis.*, vol. 28, no. 11, pp. 1479–1487, 2013.
- [3] S. Milstien *et al.*, “Sphingosine-1-phosphate is a missing cofactor for the E3 ubiquitin ligase TRAF2,” *Nature*, vol. 465, no. 7301, pp. 1084–1088, 2010.
- [4] G. Chen and D. V Goeddel, “TNF-R1 Signaling : A Beautiful Pathway The Fas Signaling Pathway,” *Science (80-.)*, vol. 296, no. May, pp. 1634–1635, 2002.
- [5] R. Medeiros *et al.*, “The role of TNF- α signaling pathway on COX-2 upregulation and cognitive decline induced by β -amyloid peptide,” *Behav. Brain Res.*, vol. 209, no. 1, pp. 165–173, 2010.
- [6] R. M. Breyer, C. K. Bagdassarian, S. A. Myers, and M. D. Breyer, “ P ROSTANOID R ECEPTORS : Subtypes and Signaling ,” *Annu. Rev. Pharmacol. Toxicol.*, vol. 41, no. 1, pp. 661–690, 2002.
- [7] D. Wang, F. G. Buchanan, H. Wang, S. K. Dey, and R. N. DuBois, “Prostaglandin E 2 enhances intestinal adenoma growth via activation of the ras-mitogen-activated protein kinase cascade,” *Cancer Res.*, vol. 65, no. 5, pp. 1822–1829, 2005.
- [8] D. Trufa, C. Übel, H. Sirbu, S. Finotto, and S. Mousset, “Establishing the role of tyrosine kinase 2 in cancer,” *Oncoimmunology*, vol. 2, no. 1, p. e22840, 2013.
- [9] A. Jain, S. Kaczanowska, and E. Davila, “IL-1 receptor-associated kinase signaling and its role in inflammation , cancer progression , and therapy resistance,” vol. 5, no. November, pp. 1–8, 2014.
- [10] D. Bai, L. Ueno, and P. Vogt, “Akt-mediated regulation of NF κ B for the oncogenicity of PI3K and Akt,” *Int. J. Cancer*, vol. 125, no. 12, pp. 2863–2870, 2009.
- [11] M. W. L. Teng *et al.*, “IL-12 and IL-23 cytokines : from discovery to targeted therapies for immune-mediated inflammatory diseases,” vol. 21, no. 7, pp. 719–729, 2015.
- [12] M. D. Castellone and H. Teramoto, “Prostaglandin E 2 Promotes Colon Cancer Cell Growth Through a G s -Axin- b -Catenin Signaling Axis,” no. December, pp. 1504–1510, 2005.
- [13] N. Tokuhira, Y. Kitagishi, M. Suzuki, A. Minami, and A. Nakanishi, “PI3K / AKT / PTEN pathway as a target for Crohn ’ s disease therapy (Review),” pp. 10–16, 2015.
- [14] H. K. H, S. A. S, and H. Akca, “PI3K / Akt / NF- κ B Signalling Pathway on NSCLC Invasion,” vol. 6, no. 4, pp. 234–238, 2016.
- [15] A. Greenhough *et al.*, “The COX-2/PGE2 pathway: Key roles in the hallmarks of cancer and adaptation to the tumour microenvironment,” *Carcinogenesis*, vol. 30, no. 3, pp. 377–386, 2009.
- [16] Y. Shi, B. E. Paluch, X. Wang, and X. Jiang, “PTEN at a glance,” *J. Cell Sci.*, vol. 125, no. 20, pp. 4687–4692, 2012.
- [17] P. Fischer and D. Hilfiker-Kleiner, “Role of gp130-mediated signalling pathways in the

- heart and its impact on potential therapeutic aspects,” *Br. J. Pharmacol.*, vol. 153, no. SUPPL. 1, pp. 414–427, 2008.
- [18] T. Bekaii-saab, “Spotlight on bevacizumab in metastatic colorectal cancer : patient selection and perspectives,” pp. 21–30, 2016.
- [19] A. A. Al-Haidari, I. Syk, and H. Thorlacius, “MiR-155-5p positively regulates CCL17-induced colon cancer cell migration by targeting RhoA,” *Oncotarget*, vol. 8, no. 9, pp. 14887–14896, 2017.
- [20] J. Tong, L. Li, B. Ballermann, and Z. Wang, “Phosphorylation and activation of RhoA by ERK in response to epidermal growth factor stimulation,” *PLoS One*, vol. 11, no. 1, pp. 1–26, 2016.
- [21] F. Marano *et al.*, “IL-13 alters mucociliary differentiation and ciliary beating of human respiratory epithelial cells,” *J. Clin. Invest.*, vol. 108, no. 12, pp. 1817–1824, 2002.
- [22] N. Gour and M. Wills-Karp, “IL-4 and IL-13 signaling in allergic airway disease,” *Cytokine*, vol. 75, no. 1, pp. 68–78, 2015.
- [23] J. Liang *et al.*, “Sphingosine-1-Phosphate Links Persistent STAT3 Activation, Chronic Intestinal Inflammation, and Development of Colitis-Associated Cancer,” *Cancer Cell*, vol. 23, no. 1, pp. 107–120, 2013.
- [24] A. L. Hart *et al.*, “Characteristics of intestinal dendritic cells in inflammatory bowel diseases,” *Gastroenterology*, vol. 129, no. 1, pp. 50–65, 2005.
- [25] M. K. Heather L, “Signaling Pathways Supporting Tumor Invasion in Head and Neck Squamous Cell Carcinoma,” *J. Clin. Exp. Pathol.*, vol. 05, no. 03, 2015.

Supplemental Table S3. Update rules and references for computational model variables.

This table shows the rules for the update of each of the elements in the model and the references that support the rules.