

Supplementary Table 1: Characteristics of men with and without serum lipid data in REDUCE

	Men with lipid data N=6,655	Men without lipid data N=1,143	p value
Age, mean (SD)	62.3 (6.0)	63.0 (6.2)	0.077
Race, n (%)			
White	557 (8)	58 (5)	<0.0001
Non-white	6,098 (92)	1,085 (95)	
Region, n (%)			
N. America	1,898 (29)	145 (13)	<0.0001
Europe	3,937 (59)	792 (69)	
Other	820 (12)	206 (18)	
BMI (kg/m²), median (IQR)	26.9 (24.8-29.4)	26.9 (24.9-29.4)	0.387
Diabetes, n (%)			
No	6,095 (92)	1,038 (91)	0.439
Yes	560 (8)	104 (9)	
Statin use, n (%)			
No	5,438 (82)	976 (85)	0.003
Yes	1,217 (18)	167 (15)	
NSAID use, n (%)			
No	4,704 (71)	883 (77)	<0.0001
Yes	1,951 (29)	260 (23)	
Alcohol use, n (%)			
None	1,732 (26)	265 (23)	0.120
Moderate	3,183 (48)	560 (49)	
Heavy	1,709 (26)	311 (27)	
Smoking status, n (%)			
Never	3,019 (45)	532 (47)	0.453
Former	2,632 (40)	430 (38)	
Current	1,004 (15)	181 (16)	
Chronic inflammation, n (%)			
Absent	1,504 (23)	243 (21)	0.316
Present	5,151 (77)	900 (79)	
Acute inflammation, n (%)			
Absent	5,650 (85)	944 (83)	0.046
Present	1,005 (15)	199 (17)	

*data for alcohol use was missing for n=31 participants