

Supplementary Table S3

The dietary characteristics between esophageal precancerosis group and the control.

Variables †	Overall (n=382)	Control (n=195)	Case (n=187)	P value
vegetables and fruits				0.182
never	1(0.3)	1 (0.5)	0 (0.0)	
seldom	14(3.7)	10 (5.1)	4 (2.1)	
often	367(96.1)	184 (94.4)	183 (97.9)	
pickled food				0.678
never	177(46.3)	92 (47.2)	85 (45.5)	
seldom	76(19.9)	41 (21.0)	35 (18.7)	
often	129(33.8)	62 (31.8)	67 (35.8)	
fried food				0.465
never	155(40.6)	85 (43.6)	70 (37.4)	
seldom	197(51.6)	95 (48.7)	102 (54.5)	
often	30(7.9)	15 (7.7)	15 (8.0)	
hot food				0.772
never	228(59.7)	113 (57.9)	115 (61.5)	
seldom	50(13.1)	27 (13.8)	23 (12.3)	
often	104(27.2)	55 (28.2)	49 (26.2)	
mildew food				0.255
never	376(98.4)	191 (97.9)	185 (98.9)	
seldom	5(1.3)	4 (2.1)	1 (0.5)	
often	1(0.3)	0 (0.0)	1 (0.5)	

† numbers of each category (%) was given and chi-square test was used.