Supplementary Table 1. The sequences of the primers used for RT-PCR analyses

Gene	Forward primer (from 5' to 3')	Reverse primer (from 5' to 3')	Ref.
PLCε	GAAAGCTGGTCTCCTATGGC	TGCCTCTAGAAGAGAACCGG	
IL-1α	TGGCCAAAGTTCCTGACTTGTTTG	CAGGTCATTTAACCAAGTGGTGCT	S1
IL-1β	ATGGCAACTGTTCCTGAACTCAACT	CAGGACAGGTATAGATTCTTTCCTTT	S2
TNF-α	AGCCCACGTCGTAGCAAACCACCAA	ACACCCATTCCCTTCACAGAGCAAT	S3
MIP-2	CTGCCGCTCCTCAGTGCTGCACTG	GCCTTGCCTTTGTTCAGTATCTTTTGG	S4
KC	GCTTGTTCAGTTTAAAGATGGTAGGC	CGTGTTGACCATACAATATGAAAGACG	S4
IL-6	TTGCCTTCTTGGGACTGATG	CTGAAGGACTCTGGCTTTGT	S4
COX-2	CACCGTACACTACATCCTGACC	TCCTCGCTTCTGATTCTGTCTTG	S5
c-fos	AGAGGAAGAGAAACGGAGAATCC	TCCGCTTGGAGTGTATCTGTCA	
RasGRP1	GCAAGACTAGAGGCCAAATCA	ATGGTGGGGTTCTCTTTTACG	
RasGRP2	CACAGCTAGTGCGCATGTTT	ATGCTGAAAAGCTGCCTCAT	
RasGRP3	ACCTAATAAGCCTGTGGTAC	CTGCTTTGTCATGGAACTTG	
RasGRP4	CATGAATCTGGGAGTGCTGA	CAGGACTTAGCAGGCTGGAG	
Rap1A	ATGCGTGAGTACAAGCTAGTAGT	AATCTACCTCGACTTGCTTTCTG	
Rap1B	CAAGGTCAGAACCTAGCAAGAC	TGACACGATGACTTTTTGCGG	
GAPDH	GTGAAGGTCGGTGTGAACGGATTT	CACAGTCTTCTGAGTGGCAGTGAT	S6
β-actin	CTACAATGAGCTGCGTGTGG	CAACGTCACACTTCATGATGG	S7

Abbreviations used here are: IL-1, interleukin-1; TNF-α, tumor necrosis factor-α; COX, cyclooxygenase; GAPDH, glyceraldehyde-3-phosphate dehydrogenase

References for Supplementary Table 1

S1. Sato Y, Ohshima T. The expression of mRNA of proinflammatory cytokines during skin wound healing in mice: a preliminary study for forensic wound age estimation (II). Int J Legal Med 2000;113:140-5.

S2. Hara T, Saito Y, Hirai T, et al., Deficiency of protein kinase Cα in mice results in impairment of epidermal hyperplasia and enhancement of tumor formation in two-stage skin carcinogenesis. Cancer Res 2005;65:7356-62.

S3. Yoshidome H, Kato A, Edwards MJ, Lentsch AB. Interleukin-10 inhibits pulmonary NF-κB activation and lung injury induced by hepatic ischemia-reperfusion. Am J

Physiol 1999;277:L919-23.

S4. Cataisson C, Pearson AJ, Torgerson S, Nedospasov SA, Yuspa SH. Protein kinase Cα-mediated chemotaxis of neutrophils requires NF-κB activity but is independent of TNFα signaling in mouse skin in vivo. J Immunol 2005;174:1686-92.

S5. Cataisson C, Joseloff E, Murillas R, et al. Activation of cutaneous protein kinase Cα induces keratinocyte apoptosis and intraepidermal inflammation by independent signaling pathways. J Immunol 2003;171:2703-13.

S6. Inoue M, Rashid MH, Fujita R, Contos JJ, Chun J, Ueda H. Initiation of neuropathic pain requires lysophosphatidic acid receptor signaling. Nat Med 2004;10:712-8.

S7. Wu D, Tadano M, Edamatsu H, et al. Neuronal lineage-specific induction of phospholipase Cε expression in the developing mouse brain. Eur J Neurosci 2003;8:1571-80.