

Legend Supplemental Table 1

* No break point has been determined; "new" for previously unpublished mutations

References Supplemental Table 1

1. Neumann HP, Pawlu C, Peczkowska M, et al. Distinct clinical features of paraganglioma syndromes associated with SDHB and SDHD gene mutations.[erratum appears in JAMA. 2004 Oct 13;292(14):1686]. JAMA 2004;292:943-51.
2. Cascon A, Cebrian A, Ruiz-Llorente S, Telleria D, Benitez J, Robledo M. SDHB mutation analysis in familial and sporadic pheochromocytoma identifies a novel mutation. J Med Genet 2002;39:E64.
3. Neumann HP, Bausch B, McWhinney SR, et al. Germ-line mutations in nonsyndromic pheochromocytoma. N Engl J Med 2002;346:1459-66.
4. Gimenez-Roqueplo AP, Favier J, Rustin P, et al. Mutations in the SDHB gene are associated with extra-adrenal and/or malignant pheochromocytomas. Cancer Res 2003;63:5615-21.
5. Bayley JP, van Minderhout I, Weiss MM, et al. Mutation analysis of SDHB and SDHC: novel germline mutations in sporadic head and neck paraganglioma and familial paraganglioma and/or pheochromocytoma. BMC Med Genet 2006;7:1.
6. Amar L, Bertherat J, Baudin E, et al. Genetic testing in pheochromocytoma or functional paraganglioma. Journal of Clinical Oncology 2005;23:8812-8.
7. Bayley JP, Devilee P, Taschner PE. The SDH mutation database: an online resource for succinate dehydrogenase sequence variants involved in pheochromocytoma, paraganglioma and mitochondrial complex II deficiency. BMC Med Genet 2005;6:39.
8. Benn DE, Gimenez-Roqueplo AP, Reilly JR, et al. Clinical presentation and penetrance of pheochromocytoma/paraganglioma syndromes. J Clin Endocrinol Metab 2006;91:827-36.

9. Lima J, Feijao T, Ferreira da Silva A, et al. High frequency of germline succinate dehydrogenase mutations in sporadic cervical paragangliomas in northern Spain: mitochondrial succinate dehydrogenase structure-function relationships and clinical-pathological correlations. *J Clin Endocrinol Metab* 2007;92:4853-64.
10. Douwes Dekker PB, Corver WE, Hogendoorn PC, van der Mey AG, Cornelisse CJ. Multiparameter DNA flow-sorting demonstrates diploidy and SDHD wild-type gene retention in the sustentacular cell compartment of head and neck paragangliomas: chief cells are the only neoplastic component. *J Pathol* 2004;202:456-62.
11. Timmers HJ, Kozupa A, Eisenhofer G, et al. Clinical presentations, biochemical phenotypes, and genotype-phenotype correlations in patients with succinate dehydrogenase subunit B-associated pheochromocytomas and paragangliomas. *J Clin Endocrinol Metab* 2007;92:779-86.
12. Braun S, Riemann K, Kupka S, et al. Active succinate dehydrogenase (SDH) and lack of SDHD mutations in sporadic paragangliomas. *Anticancer Res* 2005;25:2809-14.
13. Amar L, Baudin E, Burnichon N, et al. Succinate dehydrogenase B gene mutations predict survival in patients with malignant pheochromocytomas or paragangliomas. *J Clin Endocrinol Metab* 2007;92:3822-8.
14. Badenhop RF, Jansen JC, Fagan PA, et al. The prevalence of SDHB, SDHC, and SDHD mutations in patients with head and neck paraganglioma and association of mutations with clinical features. *J Med Genet* 2004;41:e99.
15. Schiavi F, Boedeker CC, Bausch B, et al. Predictors and prevalence of paraganglioma syndrome associated with mutations of the SDHC gene. *JAMA* 2005;294:2057-63.
16. Niemann S, Muller U. Mutations in SDHC cause autosomal dominant paraganglioma, type 3. *Nat Genet* 2000;26:268-70.
17. Pasini B, McWhinney SR, Bei T, et al. Clinical and molecular genetics of patients with the Carney-Stratakis syndrome and germline mutations of the genes coding for the

succinate dehydrogenase subunits SDHB, SDHC, and SDHD. *Eur J Hum Genet* 2008;16:79-88.

18. Niemann S, Muller U, Engelhardt D, Lohse P. Autosomal dominant malignant and catecholamine-producing paraganglioma caused by a splice donor site mutation in SDHC. *Hum Genet* 2003;113:92-4.

19. Gimm O, Armanios M, Dziema H, Neumann HP, Eng C. Somatic and occult germline mutations in SDHD, a mitochondrial complex II gene, in nonfamilial pheochromocytoma. *Cancer Res* 2000;60:6822-5.

20. Taschner PE, Jansen JC, Baysal BE, et al. Nearly all hereditary paragangliomas in the Netherlands are caused by two founder mutations in the SDHD gene. *Genes Chromosomes Cancer* 2001;31:274-81.

21. Baysal BE, Ferrell RE, Willett-Brozick JE, et al. Mutations in SDHD, a mitochondrial complex II gene, in hereditary paraganglioma. *Science* 2000;287:848-51.

22. Cascon A, Ruiz-Llorente S, Cebrian A, et al. Identification of novel SDHD mutations in patients with phaeochromocytoma and/or paraganglioma. *Eur J Hum Genet* 2002;10:457-61.

23. Velasco A, Palomar-Asenjo V, Ganan L, et al. Mutation analysis of the SDHD gene in four kindreds with familial paraganglioma: description of one novel germline mutation. *Diagn Mol Pathol* 2005;14:109-14.

24. Dannenberg H, Dinjens WN, Abbou M, et al. Frequent germ-line succinate dehydrogenase subunit D gene mutations in patients with apparently sporadic parasympathetic paraganglioma. *Clin Cancer Res* 2002;8:2061-6.

25. Badenhop RF, Cherian S, Lord RS, Baysal BE, Taschner PE, Schofield PR. Novel mutations in the SDHD gene in pedigrees with familial carotid body paraganglioma and sensorineural hearing loss. *Genes Chromosomes Cancer* 2001;31:255-63.

26. Baysal BE, Willett-Brozick JE, Lawrence EC, et al. Prevalence of SDHB, SDHC, and SDHD germline mutations in clinic patients with head and neck paragangliomas. *J Med Genet* 2002;39:178-83.
27. Milunsky JM, Maher TA, Michels VV, Milunsky A. Novel mutations and the emergence of a common mutation in the SDHD gene causing familial paraganglioma. *Am J Med Genet* 2001;100:311-4.