

	Primary Antibodies	Secondary Antibodies	Revelation
CD4-Foxp3	- Mouse anti-Foxp3 (Abcam, Cambridge, UK) 10 µg/ml - Goat anti CD4 (R&D) 10 µg/ml	- Biotinylated F(ab') ₂ donkey anti-mouse IgG (Jackson Immunoresearch) : 2 µg/ml - FITC conjugated donkey anti-goat (Jackson) 15 µg/ml	- Cy TM 3 labeled streptavidin : 3 µg/ml (Amersham)
CD4-PD-1	Mouse anti-PD-1 (produced by D Olive) 2 µg/ml - Goat anti CD4 (R&D) 10 µg/ml	- Biotinylated F(ab') ₂ donkey anti-mouse IgG (Jackson Immunoresearch) : 2 µg/ml - FITC conjugated donkey anti-goat (Jackson) 15 µg/ml	- Cy TM 3 labeled streptavidin : 3 µg/ml (Amersham)
CD8-PD-1	- Mouse anti-PD-1 (produced by D Olive) 2 µg/ml - Rabbit-anti-CD8 (Abcam) 1 µg/ml	- Biotinylated F(ab') ₂ donkey anti-mouse IgG (Jackson Immunoresearch) : 2 µg/ml - Cyan 3 conjugated donkey anti-rabbit (Jackson Immunoresearch) : 3 µg/ml	- Cy TM 2 labeled streptavidin : 3 µg/ml (Amersham)
PD-1-PD-L1	- Goat anti-PDL-1 (R&D) 15 µg/ml - Mouse anti-PD-1 (produced by D Olive) 2 µg/ml	- FITC donkey anti-goat (Jackson Immunoresearch) 15 µg/ml - Biotinylated F(ab') ₂ donkey anti-mouse IgG (Jackson Immunoresearch) : 2 µg/ml	- Cy TM 2 labeled streptavidin : 3 µg/ml (Amersham)
Foxp3-PD-1	- Goat anti-Foxp3 (Abcam) 5 µg/ml - Mouse anti-PD-1 (produced by D Olive) 2 µg/ml	Cyan3 conjugated donkey anti-goat (Jackson Immunoresearch) 2 µg/ml - Biotinylated F(ab') ₂ donkey anti-mouse IgG (Jackson Immunoresearch) : 2 µg/ml	- Cy TM 2 labeled streptavidin : 3 µg/ml (Amersham)
Tim3-PD-1	- Goat anti-Tim-3 (R&D) 1 µg/ml - Mouse anti-PD-1 (Abcam) 2 µg/ml	- Biotinylated F(ab') ₂ rabbit anti-goat IgG (Dako) : 1 µg/ml Cyan3 conjugated donkey anti-mouse (Jackson Immunoresearch) 2 µg/ml	- Cy TM 2 labeled streptavidin : 3 µg/ml (Amersham)

Table S2 : List of antibodies used for immunofluorescence staining.