

Papillomas			
	Wild-type	Transgenic	p-value
MCP-1 / CCL2	542 ± 239	255 ± 63.5	0.053
G-CSF	456 ± 320	455 ± 303	1.00
MIP-1α / CCL3	1880 ± 565	487 ± 242	0.0031
MIP-2	1850 ± 1600	749 ± 312	0.20
TNF-α	61.2 ± 39.9	21.1 ± 12.9	0.088
M-CSF	131 ± 98.6	69.4 ± 26.8	0.24
IL-16	67700 ± 41200	9230 ± 8370	0.032
TREM-1	1670 ± 1370	336 ± 465	0.095
DMBA-TPA treated skin			
	Wild-type	Transgenic	p-value
VEGF	7.25 ± 4.65	7.20 ± 2.78	0.98
IL-12 (p40)	2.37 ± 1.51	5.60 ± 2.10	0.026
IL-4	56.4 ± 42.0	61.7 ± 29.7	0.83
IL-6	0.140 ± 0.0665	0.125 ± 0.0576	0.72
IL-10	2.388 ± 0.817	1.76 ± 1.25	0.38

Suppl. Table 1. Levels of macrophage- and macrophage polarization-associated cytokines. Values are pg of cytokine per mg of total protein, N=5. MCP-1: monocyte chemotactic protein protein 1; CSF: colony stimulating factor; G-CSF: granulocyte-CSF; MIP: macrophage inflammatory protein; M-CSF: macrophage-CSF; TREM-1: triggering receptor expressed on myeloid cells 1.