

**Supplementary Table 1. Quantitation of HMGB1 secreted by cultured tumor cells, ex vivo macrophages and MDSC, in vivo grown tumors, and cultures of MDSC differentiated from bone marrow progenitor cells.**

<b>In vitro cultured tumor cells</b>	<b>ng/ml/10<sup>6</sup> cells<sup>a</sup></b>		
4T1	36.95		
CT26	16.23		
MC38	123.23		
AT3	36.95		
B78H1	12.61		

  

<b>ng/ml/10<sup>6</sup> cells<sup>b</sup></b>			
<b>Bone marrow differentiation cultures</b>	98.61		

  

<b>Ex vivo cells</b>	<b>ng/ml/10<sup>7</sup> cells<sup>a</sup></b>		
Macrophages	0.31		
LPS-treated macrophages	0.31		
MDSC	3.29		
LPS-treated MDSC	4.24		

  

<b>In vivo grown tumors</b>	<b>Tumor weight (mg)</b>	<b>Tumor diameter (mm)</b>	<b>ng/ml/1g tumor<sup>a</sup></b>
4T1	1140	13.54	85.89
CT26	1770	9.55	173.39
B78H1	840	8.84	216.86
AT3	127	3.96	0.63

<sup>a</sup> Amount of HMGB1 after 18 hrs of culture

<sup>b</sup> Amount of HMGB1 at the end of the four day incubation