

SUPPLEMENTAL TABLES

Table 1. Mean value or prevalence of characteristics across quartiles of fibrinogen and white blood cell (WBC) count in 13,414 participants, at Visit 1 (1987-89), ARIC

Characteristics	Fibrinogen (mg/dL) quartiles				WBC count ($\times 10^9$ cells/L) quartiles			
	≤ 259	260-294	295-336	≥ 337	≤ 4.8	4.9-5.8	5.9-7.0	≥ 7.1
Mean or prevalence (%)	N=3,440	N=3,435	N=3,342	N=3,197	N=3,554	N=3,413	N=3,155	N=3,292
Age (y)	52.9	53.7	54.5	54.7	53.8	53.9	54.2	53.8
Race (% Caucasian)	80.7	78.2	73.0	63.3	62.0	75.2	79.7	80.3
Sex (% Female)	49.4	52.8	57.5	62.1	57.1	52.3	50.1	47.1
BMI (kg/m ²)	26.3	27.2	28.0	29.5	27.0	27.7	28.2	28.1
Sport index	2.6	2.5	2.5	2.3	2.5	2.5	2.4	2.4
>High school education, (%)	51.6	47.4	41.7	37.3	46.2	45.9	44.3	42.0
Current smokers, (%)	14.7	21.9	29.1	35.6	10.1	15.9	24.6	50.9
Current alcohol intake, (%)	63.4	60.2	54.1	47.2	51.0	57.6	57.1	60.3
Aspirin, recent use (%)	47.5	46.0	45.6	46.2	42.5	46.3	47.4	49.5
Diabetes (%)	6.9	9.0	11.5	16.3	7.6	8.6	12.4	15.2
Current HRT, (% in females)	27.6	21.1	17.6	12.3	20.9	18.7	18.6	18.9

*Geometric means are presented for WBC since WBC was log transformed.

Table 2. Mean value or prevalence of characteristics across quartiles of von Willebrand factor (VWF) and Factor VIII (FVIII) in 13,414 participants, at Visit 1 (1987-89), ARIC

Characteristics	VWF (%) quartiles				FVIII (%) quartiles			
	≤83	84-109	110-142	≥143	≤104	105-126	127-150	≥151
Mean or prevalence (%)	N=3,430	N=3,400	N=3,365	N=3,219	N=3,347	N=3,526	N=3,265	N=3,276
Age (y)	52.5	53.8	54.4	55.1	52.7	53.7	54.3	54.9
Race (% Caucasian)	80.8	79.1	74.5	60.7	83.8	79.2	73.5	59.0
Sex (% Female)	55.1	55.0	53.8	56.5	49.8	52.9	56.2	61.7
BMI (kg/m ²)	26.9	27.3	27.9	28.9	26.5	27.3	27.8	29.2
Sport index	2.5	2.5	2.4	2.4	2.5	2.5	2.4	2.3
>High school education, (%)	49.0	46.6	44.0	38.6	50.4	46.3	44.1	37.5
Current smokers, (%)	24.5	24.6	24.9	26.0	29.5	26.3	23.0	21.0
Current alcohol intake, (%)	62.9	59.1	55.9	47.0	65.0	59.4	53.8	46.9
Aspirin, recent use (%)	47.9	48.0	45.2	44.0	47.5	46.8	46.5	44.5
Diabetes (%)	5.7	8.5	11.7	17.8	4.1	7.6	11.1	21.0
Current HRT, (% in females)	21.0	21.0	19.0	16.3	21.1	20.0	21.0	15.9

Table 3. Mean value or prevalence of characteristics across albumin quartiles in 13,414 participants, at Visit 1 (1987-89), ARIC

Characteristics	Albumin (g/dL) quartiles			
	≤3.7	3.8	3.8-3.9	>3.9
Mean or prevalence (%)	N=4,096	N=2,008	N=3,960	N=3,350
Age (y)	54.3	54.2	53.9	53.3
Race (% Caucasian)	65.1	76.1	78.1	78.8
Sex (% Female)	66.6	55.5	52.4	43.9
BMI (kg/m ²)	29.1	27.7	27.2	26.7
WHR	0.92	0.92	0.93	0.93
Sport index	2.3	2.4	2.5	2.5
>High school education, (%)	41.7	45.5	45.6	46.6
Current smokers, (%)	28.2	25.7	23.7	22.3
Current alcohol intake, (%)	49.9	56.3	58.5	61.9
Aspirin, (%)	47.5	47.0	46.4	44.5
Diabetes (%)	13.5	9.6	10.0	9.2
Current HRT, (% in females)	24.5	20.8	16.4	12.9