

Supplementary Table 3 - Baseline characteristics for women with and without missing data for biomarkers

	No missing biomarker data N=390	Missing biomarker data N=79
Endometrial Cancer Cases; n(%)	131 (34)	32 (41)
Age at blood collection, <i>years</i> ; median (interquartile range)	60.1 (56.7-63.3)	59.4 (55.8-62.1)
Age at the end of follow up, <i>years</i> ; median (interquartile range)	71.0 (65.6-76.6)	69.4 (62.1-76.1)
Fasting status at blood draw, <i>hours</i> ; n(%)		
<3	207 (53)	32 (41)
3-6	67 (17)	16 (20)
>6	116 (30)	31 (39)
Region of recruitment; n(%)		
Western Europe	110 (28)	10 (13)
Northern Europe	148 (38)	37 (47)
Southern Europe	132 (34)	32 (41)
Full-term pregnancies, <i>number</i> ; median (interquartile range)	2.0 (1.0-3.0)	2.0 (1.0-2.0)
Age at menarche, <i>years</i> ; median (interquartile range)	13.0 (12.0-14.0)	13.0 (12.0-14.0)
Had history of oral contraceptive use; n(%)	132 (34)	24 (30)
Had history of hormone therapy; n(%)	65 (17)	18 (23)
Smoking status; n(%)		
never	260 (67)	48 (61)
former	73 (19)	14 (18)
current	57 (15)	17 (22)
Physical activity; n(%)		
Inactive to moderately inactive	139 (36)	31 (39)
moderately active to active	251 (64)	48 (61)
Body Mass Index, <i>kg/m²</i> ; n(%)		
≥18.5 & <25	151 (39)	25 (32)
≥25 & <30	152 (39)	30 (38)
≥30	87 (22)	24 (30)
median (IQR)	26.3 (23.6-29.5)	27.2 (23.4-30.8)
Waist Circumference, <i>cm</i> ; n(%)		
≤80	149 (38)	27 (34)
>80 & ≤88	102 (26)	23 (29)
>88	139 (36)	29 (37)
median (interquartile range)	84.1 (77.0-92.8)	84.0 (79.0-96.0)
Waist-hip ratio; n(%)		
≤0.78	130 (33)	23 (29)
>0.78 & ≤0.84	139 (36)	27 (34)
>0.84	121 (31)	29 (37)
median (interquartile range)	0.8 (0.8-0.9)	0.8 (0.8-0.9)