

Supplementary table 1. Primary investigation series. Cox's proportional hazard regression model used to estimate the prognostic value of pStathmin(S38) in endometrial carcinomas, in relation to histopathologic variables and Stathmin.

Variable	N (%)	Unadjusted HR ^a	95% CI	P ^b	Adjusted HR	95% CI	P ^b
Histologic subtype			3.4-9.5	<0.001		1.3-4.9	0.009
Endometrioid	366 (82)	1			1		
Non-endometrioid	82 (18)	5.7			2.5		
Histologic grade			3.3-9.5	<0.001		1.1-4.7	0.03
Grade 1/2	308 (69)	1			1		
Grade 3	82 (18)	5.6			2.3		
Myometrial infiltration			3.7-12.3	<0.001		3.2-10.8	<0.001
< 50%	284 (63)	1			1		
≥ 50%	164 (37)	6.8			5.9		
Stathmin						0.7-2.4	0.4
Low	371 (83)	1	1.4-4.1	0.002	1		
High	77 (17)	2.4			1.3		
pStathmin(S38)			1.6-4.4	<0.001		1.0-3.1	0.05
Low	326 (73)	1			1		
High	122 (27)	2.6			1.8		

N: number of cases; HR: Hazard ratio; CI: confidence interval; FIGO: International Federation of Gynecology and Obstetrics

^aUnadjusted HRs given for analyses of 448 cases with data available for all variables in the multivariate analysis; ^bLikelihood ratio test