


Supplementary Figure 3.



B

Steady State Pharmacokinetic Parameters for Dose Level -1A Olaparib 100mg BID and Prexasertib 70 mg/m² IV

Parameter	Olaparib 100 mg BID Alone	Olaparib 100mg BID + Prexasertib 70mg/m ²	Ratio*	<i>P</i> value
Median Range <i>T</i> _{max} (h)	1.95 (0.5-4.0)	2.98 (1.0-4.0)	1.53	0.20
<i>C</i> _{max} (ng/mL)	6,3332 (33.0)	5,260 (35.7)	0.83	0.26
<i>C</i> _{min} (ng/mL)	1002 (90.3)	726 (85.5)	0.72	0.35
AUC _τ (ng.h/mL)	33037 (44.9)	27587 (43.6)	0.84	0.38

Values are reported as the geometric mean %CV) for all other parameters.
*Mean parameter when combined with prexasertib divided by mean parameter for olaparib alone during cycle 0

Figure S3.

A. Prexasertib Plasma Levels.

Plasma concentrations (ng/mL) at 70mg/m² and 80mg/m² doses are shown. Mean prexasertib exposures for patients enrolled to the 70 mg/m² and 80mg/m² dose levels were 717.4 ng/mL (range 310-1070 ng/mL) and 721.1 ng/mL (range from 327-1090 ng/mL) 60 minutes after dosing.

B. Pharmacokinetic Parameters for Olaparib Alone and in Combination with Prexasertib.

Shown are the pharmacokinetic parameters for Dose Level -1A: Olaparib 100mg PO BID administered alone and in combination with prexasertib 70mg/m².