

**Supplementary Table S5.** Incidence of lymphopenia and lymphocyte decreased, regardless of attribution

MedDRA preferred term, <i>n</i> (%)	0.2 mg ( <i>n</i> =3)		0.8 mg ( <i>n</i> =3)		3.2 mg ( <i>n</i> =6)		12 mg ( <i>n</i> =11)		40 mg ( <i>n</i> =7)		80 mg ( <i>n</i> =12)		160 mg ( <i>n</i> =11)		300 mg ( <i>n</i> =109)		600 mg ( <i>n</i> =6)		1200 mg ( <i>n</i> =4)		All Patients ( <i>N</i> =172)			
	Grade		Grade		Grade		Grade		Grade		Grade		Grade		Grade		Grade		Grade		Grade			
	All	≥3	All	≥3	All	≥3	All	≥3	All	≥3	All	≥3	All	≥3	All	≥3	All	≥3	All	≥3	All	≥3		
<b>Lymphopenia</b>	0	0	0	0	1 (17)	1 (17)	1 (9)	1 (9)	1 (14)	1 (14)	2 (17)	2 (17)	1 (9)	1 (9)	5 (5)	4 (4)	0	0	0	0	11 (6)	10 (6)		
<b>Lymphocyte count decreased</b>	1 (33)	1 (33)	0	0	0	0	1 (9)	1 (9)	0	0	0	0	0	0	1 (1)	0	0	0	0	0	0	0	3 (2)	2 (1)