

## Supplementary Table S2

**Supplementary Table S2.** Immunohistochemistry of mouse tumors treated for two weeks with vehicle control, niclosamide (20 mg/kg), TRA-8 (200  $\mu$ g) or the combination of both agents. Tumors were stained and analyzed (H-score) for p-STAT3, cytosolic  $\beta$ -catenin, nuclear  $\beta$ -catenin and cleaved caspase 3. The values represent an average of three tumors per group ( $\pm$  SE).

	p-STAT3	$\beta$ -Catenin (cytoplasm)	$\beta$ -Catenin (nucleus)	Cleaved caspase- 3
<b>Vehicle Control</b>	55.0 ( $\pm$ 28)	76.7 ( $\pm$ 37)	20.0 ( $\pm$ 12)	45.0 ( $\pm$ 38)
<b>Niclosamide (20 mg/kg)</b>	0.0 ( $\pm$ 0)	53.3 ( $\pm$ 13)	6.7 ( $\pm$ 7)	40.0 ( $\pm$ 30)
<b>TRA-8 (200 <math>\mu</math>g)</b>	13.3 ( $\pm$ 7)	50.0 ( $\pm$ 21)	0.0 ( $\pm$ 0)	76.7 ( $\pm$ 38)
<b>Niclosamide and TRA-8</b>	0.0 ( $\pm$ 0)	23.3 ( $\pm$ 13)	6.7 ( $\pm$ 3)	85.0 ( $\pm$ 43)