

Supplementary Table 1. Clinicopathologic factors of patients analyzed b
miRNA array

Clinicopathological variables	Non-liver metastasis cases (n = 12)	Liver metastasis cases (n = 7)	P value
Gender			
Male	6 (50.0%)	4 (57.1%)	
Female	6 (50.0%)	3 (42.9%)	NS
Age (mean \pm SD)	56.8 \pm 11.8	64.3 \pm 8.38	NS
CRC tumor size (mean \pm SD)	3.78 \pm 2.31	4.27 \pm 2.14	NS
Histological grade ^a			
Well/mod	12 (100%)	6 (85.7%)	
Others	0 (0%)	1 (14.3%)	NS
Tumor invasion			
Tis/T1	0 (0%)	0 (0%)	
T2	7 (58.7%)	1 (14.3%)	
T3	4 (33.3%)	6 (85.7%)	
T4	1 (8.33%)	0 (0%)	NS
UICC Stage ^b			
I	6 (50%)	0 (0%)	
II	4 (33.3%)	0 (0%)	
III	2 (16.7%)	2 (28.6%)	
IV	0 (0%)	5 (71.4%)	< 0.001 ^c
Lymphatic invasion			
Absent	4 (33.3%)	0 (0%)	
Present	8 (66.7%)	7 (100%)	NS
Venous invasion			
Absent	10 (83.3%)	3 (42.9%)	
Present	2 (16.7%)	4 (57.1%)	NS
Lymph node metastasis			
Absent	10 (83.3%)	0 (0%)	
Present	2 (16.7%)	7 (100%)	< 0.001 ^c

^aHistological grade. Well, well differentiated adenocarcinoma; Mod, moderately differentiated adenocacinoma; Others, poorly differentiated and mucinous adenocaricinoma.